This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Original) Liquid-crystalline medium based on a mixture of polar compounds of positive dielectric anisotropy, characterised in that it comprises one or more compounds of the formula I

$$R^1 \longrightarrow H \longrightarrow L^1$$
 L^2
 I

and one or more compounds of the formula IA

$$R^2$$
 H A Z^1 B Z^2 O X^2 IA

where the proportion of the compounds of the formula I in the medium is at least 18% by weight, and in which the individual radicals have the following meanings:

- R¹ is an alkenyl radical having from 2 to 8 carbon atoms,
- is H, an alkyl radical having from 1 to 15 carbon atoms which is halogenated, substituted by CN or CF_3 or unsubstituted, where, in addition, one or more CH_2 groups in these radicals may each, independently of one another, be replaced by $-C \equiv C_7$, $-CO_7$, $-CH = CH_7$, $-O_7$, $-CH_7$ in

such a way that O atoms are not linked directly to one another,

X¹ is an alkyl radical, alkenyl radical, alkoxy radical or alkenyloxy radical, each having up to 6 carbon atoms, in the case where a =

1 also F, Cl, CN, SF₅, SCN, NCS or OCN,

is F, Cl, CN, SF₅, SCN, NCS, OCN, a halogenated alkyl radical, halogenated alkenyl radical, halogenated alkoxy radical or halogenated alkenyloxy radical, each having up to 6 carbon atoms,

 Z^1 and Z^2 are each, independently of one another, -CF₂O-, -OCF₂- or a single bond, where $Z^1 \neq Z^2$,


a is 0 or 1, and

 L^{14} are each, independently of one another, H or F.

2. (Original) Liquid-crystalline medium according to Claim 1, characterised in that it comprises one, two or more compounds of the formulae IA-1 to IA-30

$$R^2$$
 H O O CF_2O O F IA-1


$$R^2$$
 H O O CF_2O O F $IA-2$



$$R^2$$
 H O O CF_2O O O $IA-4$

$$R^2$$
 H O CF₂O O OCHF₂ IA-7

$$R^2$$
 H O CF₂O O CHF₂ IA-8



$$R^2$$
 H O CF₂O O CI IA-10

$$R^2$$
 H O CF₂O O CI IA-11

$$R^2$$
 H O O CF_2O O F IA-12

$$R^2$$
 H O CF_2O O CF_3 IA-13

$$R^2$$
 H O CF_2O O CF_3 IA-14

$$R^2$$
 H O O CF_2O O F CF_3 $IA-15$

$$R^2$$
 H O CF_2O O F IA-16

$$R^2$$
 H O CF_2O O F IA-17

$$R^2$$
 H O F CF_2O F IA-18

$$R^2$$
 H O CF_2O O O $IA-19$

$$R^2$$
 H O CF_2O O O $IA-22$

$$R^2$$
 H O CF_2O O $COHF_2$ IA-23

$$R^2$$
 H O CF₂O O CI IA-25

$$R^2$$
 H O CF₂O O CI IA-26

$$R^2$$
 H O F CF_2O O F $IA-27$

$$R^2$$
 H O CF_2O O CF_3 IA-28

$$R^2$$
 H O CF_2O O CF_3 $IA-29$

$$R^2$$
 H O CF_2O O CF_3 $IA-30$

in which R² is as defined in Claim 1.

3. (Currently Amended) Liquid-crystalline medium according to Claim 1 or 2, characterised in that it comprises one or more compounds of the formulae I-1 to I-5

alkenyl—
$$H$$
— O — F

alkenyl—
$$H$$
— O — F

$$(F)$$

in which alkenyl is an alkenyl radical having from 2 to 8 carbon atoms and alkyl is a straight-chain alkyl radical having 1-15 carbon atoms.

4. (Currently Amended) Liquid-crystalline medium according to one of Claims 1 to 3 Claim 1, characterised in that it additionally comprises one or more compounds selected from the group consisting of the general formulae II, III, IV, V and VI

$$R^0$$
 H
 O
 X^0
II

$$R^0 \longrightarrow H$$
 $C_2H_4 \longrightarrow O$ $C_2H_4 \longrightarrow O$ III

$$R^0$$
 H O Y^3 Y^1 V Y^0 V

- 9 -

$$R^{0} \longrightarrow H \longrightarrow Z^{0} \longrightarrow H \longrightarrow Q^{1} \longrightarrow X^{0} \longrightarrow Y^{1}$$

$$R^{0} \longrightarrow H \longrightarrow Z^{0} \longrightarrow Q^{1} \longrightarrow Y^{1} \longrightarrow$$

in which the individual radicals have the following meanings:

and the compound of the formula II is not identical with the compound of the formula I.

- 5. (Original) Liquid-crystalline medium according to Claim 4, characterised in that the proportion of compounds of the formulae IA and I to VI together in the mixture as a whole is at least 50% by weight.
- 6. (Currently Amended) Liquid-crystalline medium according to one of Claims 1 to 5 Claim 1, characterised in that it additionally comprises one or more compounds of the formulae Ea to Ef

$$R^0$$
 H H COO O F Ea

$$R^0$$
 H COO O F Ec

$$R^0$$
 H H COO O OCF_3 Eb

$$R^0$$
 H COO O F OCF_3 Ed


$$R^0$$
 H O COO O F Ee

$$^{\circ}R^{0}$$
 \longrightarrow $^{\circ}COO$ \longrightarrow O \longrightarrow COO \longrightarrow OCF_{3}

in which R⁰ is as defined in Claim 4 H, n-alkyl, alkoxy, oxaalkyl, fluoroalkyl, alkenyloxy or alkenyl, each having up to 9 carbon atoms.

7. (Currently Amended) Liquid-crystalline medium according to one of Claims 1 to 6 Claim 1, characterised in that it comprises one or more compounds of the

formulae IIa to IIg



$$R^0$$
 H O F IIb

$$R^0$$
 H O OCHFCF₃

$$R^0$$
 H O OCHFCF₃ IId

$$R^0$$
 H O O OCHFCF₃ IIe

$$R^0$$
 H O OCHF₂ IIf

$$R^0$$
 H O CHF₂ IIg

in which R⁰ is as defined in Claim 4 is H, n-alkyl, alkoxy, oxaalkyl, fluoroalkyl, alkenyloxy or alkenyl, each having up to 9 carbon atoms.

8. (Currently Amended) Liquid-crystalline medium according to one of Claims 1 to 7 Claim 1, characterised in that it additionally comprises one or more compounds of the formulae RI to RVII

$$R^* - H - CF_3$$
 RII

$$R^*$$
 H OCF_3 RIII

$$R^* - H - (O)CH = CF_2$$
 RVI

$$R^* - H - (O)CF = CF_2$$
 RVII

in which

R* is n-alkyl, alkoxy, oxaalkyl, fluoroalkyl or alkenyloxy, each having up to 9 carbon atoms, and

alkyl and

alkyl* are each, independently of one another, a straight-chain or branched

alkyl radical having 1-9 carbon atoms.

- 9. (Currently Amended) Liquid-crystalline medium according to one of Claims 1 to 8 Claim 1, characterised in that the proportion of compounds of the formula IA in the mixture as a whole is from 5 to 40% by weight.
- 10. (Original) Use of the liquid-crystalline medium according to Claim 1 for electro-optical purposes.
- 11. (Original) Electro-optical liquid-crystal display containing a liquid-crystalline medium according to Claim 1.

DOCKET NO.: MERCK-3119